

Intervento	Antibiotico di prima scelta	Alternativa nei pazienti allergici ai betalattamici	Note
Parto cesareo elettivo	Cefazolina 2 g ev (dopo clampaggio del cordone ombelicale)	Clindamicina 600 mg ev (dopo clampaggio del cordone ombelicale)	
Parto cesareo non elettivo (con travaglio in atto e/o rottura di membrane più di 6 ore prima dell'intervento)	Cefoxitina 2 gr (dopo clampaggio del cordone ombelicale)	Clindamicina 600 mg ev +/-Gentamicina 3 mg/kg ev (dopo clampaggio cordone ombelicale)	
Isterectomia addominale e vaginale	Cefoxitina 2 gr	Clindamicina 600 mg +/-Gentamicina 3 mg/kg ev	
Aborto indotto	Doxiciclina 200 mg per os 2 ore prima dell'intervento	Claritromicina 500 mg per os 2 ore prima dell'intervento	

CHIRURGIA ORTOPEDICA [84, 87, 369-392]

Intervento	Antibiotico di prima scelta	Alternativa nei pazienti allergici ai betalattamici	Note
Chirurgia elettiva senza protesi (meniscectomia e sinoviectomia artroscopica, riparazione di muscoli, tendini e fasce)	No		
Chirurgia osteoarticolare, ricostruzione di LCA ginocchio, sinoviectomia con artrotomia, artrodesi)	Cefazolina 2 g ev	Clindamicina 600 mg ev	
Artroprotesi (anca, ginocchio altre protesi)	Cefazolina 2 g ev	Vancomicina 1 g ev	
Fissazione di frattura chiusa	Cefazolina 2 g ev	Clindamicina 600 mg ev	

Chirurgia del rachide	Cefazolina 2 g ev	Vancomicina 15 mg/Kg ev (max 1 g)
-----------------------	-------------------	--------------------------------------

12. BIBLIOGRAFIA

- 1- Bernard HR, Cole WR. The prophylaxis of surgical infection: the effect of prophylactic antimicrobial drugs on the incidence of infection following potentially contaminated operations. *Surgery* 1964;56:151-7
- 2- Polk HC Jr, Lopez-Mayor JF. Postoperative wound infection: a prospective study of determinant factors and Prevention. *Surgery* 1969; 66(1): 97-103
- 3- Stone HH. Basic principles in the use of prophylactic antibiotics. *J Antimicrob Chemother.* 1984; 14 Suppl B: 33-7
- 4- Meijer WS, Schmitz PI, Jeekel J. Meta-analysis of randomized, controlled clinical trials of antibiotic prophylaxis in biliary tract surgery. *Br J Surg.* 1990; 77 (3): 283-90
- 5- Roach A, Kernodle D, Kaiser A. Selecting cost-effective antimicrobial prophylaxis in surgery: are we getting what we pay for? *DICPAnn Pharmacother.* 1990; 24: 183-5
- 6- Aznar R, Mateu M, Mirò JM, Gatell JM, Gimferrer JM, Aznar E, Mallolas J, Soriano E, Sanchez-Lloret J. Antibiotic prophylaxis in non-cardiac thoracic surgery: cefazolin versus placebo. *Eur J Cardiothorac Surg.* 1991; 5 (10): 515-8
- 7- McGowan JE Jr. Cost and benefit of perioperative antimicrobial prophylaxis: methods for economic analysis. *Rev Infect Dis.* 1991 ; 13 Suppl 10: S879-89
- 8- Davey P, Malek M, Thomas P. Measuring the cost-effectiveness of antibiotic prophylaxis in surgery. *Am J Surg.* 1992; 164 (4A Suppl): 39S-43S
- 9- Page CP, Bohnen JMA, Fletcher JR, et al. Antimicrobial prophylaxis for surgical wounds: guidelines for clinical care. *Arch Surg.* 1993; 128: 79-88
- 10- Dellinger EP, Gross PA, Barrett TL, Krause PJ, Martone WJ, McGowan JE Jr, Sweet RL, Wenzel RP. Quality standard for antimicrobial prophylaxis in surgical procedures. Infections Diseases Society of America. *Clin Infect Dis.* 1994; 18 (3): 422-7
- 11- Lizàn-Garcia M, Garcia-Caballero J, Asensio-Vegas A. Risk factors for surgical-wound infection in general surgery: a prospective study. *Infect Control Hosp Epidemiol.* 1997; 18 (5): 310-5
- 12- Sasse A, Mertens R, Sion JP, Ronveaux O, Bossens M, De Mol P, Goossens H, Lauwers S, Potvliege C, Van Landuyt H, Verbist L, Verschraegen G. Surgical prophylaxis in Belgian hospitals: estimate of costs and potential savings. *J Antimicrob Chemother.* 1998; 41 (2): 267-72
- 13- Külling D, Sonnenberg A, Fried M, Bauerfeind P. Cost analysis of antibiotic prophylaxis for PEG. *Gastrointest Endosc.* 2000; 51 (2): 152-6

- 14- Fernández AH, Monge V, Garcinuno MA. Surgical antibiotic prophylaxis: effect in postoperative infections *Eur J Epidemiol.* 2001; 17(4): 369-74
- 15- Hartmann B, Sucke J, Brammen D, Jost A, Eicher A, Róhrig R, Junger A. Impact of inadequate surgical antibiotic prophylaxis on perioperative outcome and length of stay on ICU in general and trauma surgery. Analysis using automated data collection. *Int J Antimicrob Agents.* 2005; 25 (3): 231-6
- 16- Silber JH, Rosenbaum PR, Trudeau ME, Chen W, Zhang X, Lorch SA, Kelz RR, Mosher RE, Even-Shoshan O. Preoperative antibiotics and mortality in the elderly. *Ann Surg.* 2005; 242 (1): 107-14
- 17- Manniën J, van Kasteren ME, Nagelkerke NJ, Gyssens IC, Kullberg BJ, Wille JC, de Boer AS. Effect of optimized antibiotic prophylaxis on the incidence of surgical site infection. *Infect Control Hosp Epidemiol.* 2006; 27 (12): 1340-6
- 18- Agenzia Regionale della Sanità Regione autonoma Friuli-Venezia Giulia. Linee guida per la profilassi antibiotica in chirurgia. (<http://www.vvvy.sanita.fvg.it/ars/specializza/progetti/allegati/tot-00.pdn>)
- 19- Regione Emilia-Romagna - Agenzia sanitaria regionale. Linee guida per la chemioprolifassi antibiotica in chirurgia. Indagine conoscitiva in Emilia-Romagna. Dossier 72 2002. (http://asr.regione.emilia-romagna.it/wcm/asr/collana_dossier/doss072/link/doss72.pdf)
- 20- American Society of Health-System Pharmacists. ASHP Therapeutic Guidelines on Antimicrobial Prophylaxis in Surgery, *AM J Health-Syst-Pharm.* 1999; 56: 1839-88
- 21- Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR Guideline for prevention of surgical site infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control.* 1999; 27: 97-132
- 22- SIGN - Scottish Intercollegiate Guidelines Network. Antibiotic prophylaxis in surgery - A national clinical guideline. SIGN publication n. 104, Edinburgh, July 2008) (<http://www.sign.ac.uk/pdf/sign104.pdf>)
- 23- Sistema Nazionale Linee Guida. Documento 17 - Antibiotico-profilassi perioperatoria nell'adulto. Settembre 2008 (http://www.snlq-iss.it/cms/files/LG_Antibioticoprofilassi_Unico_2008.pdf)
- 24- Bratzler DW, Houck PM, Surgical Infection Prevention Guidelines Writers Workgroup et al. Antimicrobial prophylaxis for surgery: an advisory statement from the National Surgical Infection Prevention Project. *Clin Infect Dis.* 2004; 38 (12): 1706-15
- 25- No authors listed. Antimicrobial prophylaxis for surgery, *Treat Guidel Med Leti.* 2006; 4 (52): 83-8

- 26- ICSI - Institute for Clinical Systems Improvement. Health Care Protocol: Prevention of Surgical Site Infection. 2006 (http://www.icsi.org/surgical_site_infection_protocol/surgical_site_infection_prevention_protocol_pdf.html)
- 27- Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG. CDC definitions of nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound infections. *Infect Control Hosp Epidemiol.* 1992; 13 (10): 606-8
- 28- Green JW, Wenzel RP. Postoperative wound infection: a controlled study of the increased duration of hospital stay and direct cost of hospitalization. *Ann Surg.* 1977;185 (3): 264-8
- 29- Taylor GJ, Mikell FL, Moses HW, Dove JT, Katholi RE, Malik SA, Markwell SJ, Kormeyer C, Schneider JA, Wellons HA. Determinants of hospital charges for coronary artery bypass surgery: the economic consequences of postoperative complications. *Am J Cardiol.* 1990; 65 (5): 309-13
- 30- Wenzel RP. Preoperative antibiotic prophylaxis. *N Engl J Med* 1992; 326(5): 337-9
- 31- Kirkland KB, Briggs JP, Trivette SL, Wilkinson WE, Sexton DJ The impact of surgical site infections in the 1990s: attributable mortality, excess length of hospitalization, and extra costs. *Infect Control Hosp Epidemiol.* 1999; 20 (11): 725-30
- 32- Hollenbeak CS, Murphy D, Dunagan WC, Fraser VJ. Nonrandom selection and the attributable cost of surgical-site infections. *Infect Control Hosp Epidemiol.* 2002; 23 (4): 177-182
- 33- Whitehouse JD, Friedman ND, Kirkland KB, Richardson WJ, Sexton DJ. The impact of surgical site-infections following orthopedic surgery at a community hospital and a university hospital: adverse quality of life, excess length of stay, and extra cost. *Infect Control Hosp Epidemiol.* 2002; 23 (4): 183-9
- 34- Fry DE. The economic costs of surgical site infection. *Surg Infect (Larchmt).* 2002; 3 Suppl 1 : S37-43
- 35- Perencevich EN, Sands KE, Cosgrove SE, Guadagnoli E, Meara E, Platt R. Health and economic impact of surgical site infections diagnosed after hospital discharge. *Emerg Infect Dis.* 2003; 9 (2): 196-203
- 36- McGarry SA, Engemann JJ, Schmader K, Sexton DJ, Kaye KS. Surgical-site infection due to *Staphylococcus aureus* among elderly patients: mortality, duration of hospitalization, and cost. *Infect Control Hosp Epidemiol.* 2004; 25 (6): 461-7
- 37- Leaper DJ, van Goor H, Reilly J, Petrosillo N, Geiss HK, Torres AJ, Berger A. Surgical site infection - a European perspective of incidence and economic burden. *Int Wound J.* 2004; 1 (4): 247-73
- 38- Urban JA. Cost analysis of surgical site infections. *Surg Infect (Larchmt).* 2006; 7 Suppl 1: S19-22

- 39- Penel N, Lefebvre JL, Cazin JL, Clisant S, Neu JC, Dervaux B, Yazdanpanah Y. Additional direct medical costs associated with nosocomial infections after head and neck cancer surgery: a hospital-perspective analysis. *Int J Oral Maxillofac Surg.* 2008; 37 (2): 135-9
- 40- Fukatsu K, Saito H, Matsuda T, Ikeda S, Furukawa S, Muto T. Influences of type and duration of antimicrobial prophylaxis on an outbreak of methicillin-resistant *Staphylococcus aureus* and on the incidence of wound infection. *Arch Surg.* 1997; 132 (12): 1320-5
- 41- Shlaes DM, Gerding DN, John JF Jr, Craig WA, Bornstein DL, Duncan RA, Eckman MR, Farrer WE, Greene WH, Lorian V, Levy S, McGowan JE Jr, Paul SM, Ruskin J, Tenover FC, Watanakunakorn C. Society for Healthcare Epidemiology of America and Infectious Diseases Society of America Joint Committee on the Prevention of Antimicrobial Resistance: guidelines for the prevention of antimicrobial resistance in hospitals. *Clin Infect Dis.* 1997; 25 (3): 584-99
- 42- Terpstra S, Noordhoek GT, Voesten HG, Hendriks B, Degener JE. Rapid emergence of resistant coagulase-negative staphylococci on the skin after antibiotic prophylaxis. *J Hosp Infect.* 1999; 43 (3): 195-202
- 43- Harbarth S, Samore MH, Lichtenberg D, Carmeli Y. Prolonged antibiotic prophylaxis after cardiovascular surgery and its effect on surgical site infections and antimicrobial resistance. *Circulation.* 2000; 101 (25): 2916-21
- 44- Tacconelli E, De Angelis G, Cataldo MA, Pozzi E, Cauda R. Does antibiotic exposure increase the risk of methicillin-resistant *Staphylococcus aureus* (MRSA) isolation? A systematic review and meta-analysis. *J Antimicrob Chemother.* 2008; 61 (1): 26-38
- 45- American Society for Microbiology. Report of the ASM Task Force on antimicrobial resistance. Washington: The Society, 1994
- 46- Gold HS, Moellering RC Jr. Antimicrobial-drug resistance. *N Engl J Med.* 1996; 335 (19): 1445-53
- 47- National Nosocomial Infections Surveillance System. National Nosocomial Infections Surveillance (NNIS) System Report, data summary from January 1992 through June 2004, issued October 2004. *Am J Infect Control.* 2004; 32 (8): 470-85
- 48- Privitera G, Scarpellini P, Ortisi G, Nicastro G, Nicolini R, de Lalla F. Prospective study of *Clostridium difficile* intestinal colonisation and disease following single dose antibiotic prophylaxis in surgery. *Antimicrob Agents Chemother.* 1991; 35 (1): 208-10
- 49- Kreisel D, Savel TG, Silver AL, Cunningham JD. Surgical antibiotic prophylaxis and *Clostridium difficile* toxin positivity. *Arch Surg.* 1995; 130 (9): 989-93

- 50- Spencer RC. The role of antimicrobial agents in the aetiology of Clostridium difficile-associated disease. *J Antimicrob Chemother.* 1998; 41 Suppl. C: 21-7
- 51- Harbarth S, Samore MH, Carmeli Y. Antibiotic prophylaxis and the risk of Clostridium difficile-associated diarrhoea. *J Hosp Infect.* 2001 ; 48 (2): 93-7
- 52- Wiström J, Norrby SR, Myhre EB, Eriksson S, Granström G, Lagergren L, Englund G, Nord CE, Svenungsson B. Frequency of antibiotic-associated diarrhoea in 2462 antibiotic-treated hospitalized patients: a prospective study. *J Antimicrob Chemother.* 2001; 47 (1): 43-50
- 53- Mukhtar S, Shaker H, Basarab A, Byrne JP. Prophylactic antibiotics and Clostridium difficile infection. *J Hosp Infect.* 2006; 64(1): 93-4
- 54- Carignan A, Allard C, Pépin J, Cossette B, Nault V, Valiquette L. Risk of Clostridium difficile infection after perioperative antibacterial prophylaxis before and during an outbreak of infection due to a hypervirulent strain. *Clin Infect Dis.* 2008; 46 (12): 1838-43
- 55- Cruse PJ, Foord R. A five year prospective study of 23,649 surgical wounds. *Arch. Surg.* 1973; 107 (2): 206-10
- 56- Cruse PJ, Foord R. The epidemiology of wound infection: a 10-year prospective study of 62,939 wounds. *Surg Clin North Am.* 1980; 60(1): 27-40
- 57- Haley RW, Culver DH, White JW, Morgan WM, Emori TG, Munn VP, Hooton TM. The efficacy of infection surveillance and control programs in preventing nosocomial infections in US hospitals. *Am J Epidemiol.* 1985; 121 (2): 182-205
- 58- Haley RW, Culver DH, Morgan WM, White JW, Emori TG, Hooton TM. Identifying patients at high risk of surgical wound infection. A simple multivariate index of patient susceptibility and wound contamination. *Am J Epidemiol.* 1985; 121 (2): 206-15
- 59- Culver DH, Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG, Banerjee SN, Edwards JR, Tolson JS, Henderson TS, et al. Surgical wound infection rates by wound class, operative procedure and patient risk index. National Nosocomial Infections Surveillance System. *Am J Med.* 1991; 91 (3B): 152S-157S
- 60- Garibaldi RA, Cushing D, Lerer T. Risk factors for postoperative infection. *Am J Med.* 1991; 91 (3B): 158S-163S
- 61- Society for Hospital Epidemiology of America, Robert J. Sherertz, Richard A. Garibaldi, Randy D. Marosok, C. Glen Mayhall, William E. Scheckler Association for Practitioners in Infection Control, Inc., Rosemary Berg, Centers for Disease Control, Robert P. Gaynes, William R. Jarvis, William J. Martone, Surgical Infection Society, James T. Lee Jr. Consensus paper on the

- surveillance of surgical wound infections. *Am J Infect Control*. 1992; 20(5): 263-70
- 62- Leaper DJ. Risk factors for surgical infection. *J Hosp Infect*. 1995; 30 Suppl: 127-39
- 63- Perl TM, Roy MC. Postoperative wound infections: risk factors and role of *Staphylococcus aureus* nasal carriage. *J. Chemother*. 1995; 7 Suppl 3: 29-35
- 64- de Boer AS, Mintjes-de Groot AJ, Severijnen AJ, van den Berg JM, van Pelt W. Risk assessment for surgical-site infections in orthopedic patients. *Infect Control Hosp Epidemiol*. 1999; 20 (6): 402-7
- 65- Geubbels EL, Mintjes-de Groot AJ, van den Berg JM, de Boer AS. An operating surveillance system of surgical-site infections in The Netherlands: results of the PREZIES national surveillance network. *Preventie van Ziekenhuisinfecties door Surveillance Infect Control Hosp Epidemiol*. 2000; 21 (5): 311-8
- 66- Platell C, Hall JC. The prevention of wound infection in patients undergoing colorectal surgery. *J. Hosp. Infect*. 2001; 49 (4): 233-8
- 67- Scott JD, Forrest A, Feuerstein S, Fitzpatrick P, Schentag JJ. Factors associated with postoperative infection. *Infect Control Hosp Epidemiol*. 2001; 22 (6): 347-51
- 68- Raja'a YA, Salam AR, Salih YA, Salman MS, Al-Baseer LS, Al-Kirshi NA, Al-Jalal NS. Rate and risk factors of surgical site infections with antibiotic prophylaxis. *Saudi Med J*. 2002; 23 (6): 672-4
- 69- Vazquez-Aragon P, Lizan-Garcia M, Cascales-Sanchez P, Villar-Canovas MT, Garcia-Olmo D. Nosocomial infection and related risk factors in a general surgery service: a prospective study. *J Infect*. 2003; 46 (1): 17-22
- 70- Kasatpibal N, Jamulitrat S, Chongsuvivatwong V. Standardized incidence rates of surgical site infection: a multicenter study in Thailand. *Am J Infect Control*. 2005; 33 (10): 587-94
- 71- Fiorio M, Marvaso A, Vigano F, Marchetti F. Incidence of surgical site infections in general surgery in Italy. *Infection*. 2006; 34 (6): 310-4
- 72- Cheadle WG. Risk factors for surgical site infection. *Surg Infect (Larchmt)*. 2006; 7 Suppl 1 : S7-11
- 73- Neumayer L, Hosokawa P, Itani K, El-Tamer M, Henderson WG, Khuri SF. Multivariable predictors of postoperative surgical site infection after general and vascular surgery: results from the patient safety in surgery study. *J Am Coli Surg*. 2007; 204 (6): 1178-87
- 74- American Society of Anesthesiologists. New classification of physical status. *Anesthesiology*. 1963; 24: 111

- 75- Woodfield JC, Beshay NM, Pettigrew RA, Plank LD, van Rij AM. American Society of Anesthesiologists classification of physical status as a predictor of wound infection. *ANZ J Surg.* 2007; 77 (9): 738-41
- 76- Burke JF. The effective period of preventive antibiotic action in experimental incisions and dermal lesions. *Surgery.* 1961; 50: 161-8
- 77- Stone HH, Hooper CA, Kolb LD, Geheber CE, Dawkins EJ. Antibiotic prophylaxis in gastric, biliary and colonic surgery. *Ann Surg.* 1976; 184(4): 443-52
- 78- Galandiuk S, Polk HC Jr, Jagelman DG, Fazio VW. Re-emphasis of priorities in surgical antibiotic prophylaxis. *Surg Gynecol Obstet.* 1989; 169 (3): 219-22
- 79- Classen DC, Evans RS, Pestotnik SL, Horn SD, Menlove RL, Burke JP. The timing of prophylactic administration of antibiotics and the risk of surgical- wound infection. *N Engl J Med.* 1992; 326 (5): 281-6
- 80- Wong-Beringer A, Corelli RL, Schrock TR, Guglielmo BJ. Influence of timing of antibiotic administration on tissue concentrations during surgery. *Am J Surg.* 1995; 169 (4): 379-81
- 81- Kernodle DS, Kaiser AB. Surgical and trauma-related infections. In: Mandell GL, Bennet JE, Dolin R, eds. *Principles and Practice of Infectious Diseases*, 4th edn. New York: Churchill Livingstone, 1995; 2742-55
- 82- Woods RK, Dellinger EP. Current guidelines for antibiotic prophylaxis of surgical wounds. *Am Fam Physician.* 1998; 57 (11): 2731-40
- 83- Gyssens IC. Preventing postoperative infections: current treatment recommendations. *Drugs.* 1999; 57 (2): 175-85
- 84- American Academy of Orthopaedics (AAOS). Advisory Statement. Recommendations for the use of intravenous antibiotic prophylaxis in primary total joint arthroplasty. 2004. (<http://www.aaos.org/about/papers/advistmt/1027.asp>)
- 85- Bratzler DW, Houck PM, Richards C, Steele L, Dellinger EP, Fry DE, Wright C, Ma A, Carr K, Red L. Use of antimicrobial prophylaxis for major surgery: baseline results from the National Surgical Infection Prevention Project. *Arch Surg.* 2005; 140 (2): 174-82
- 86- Gilbert DN, Moellering RC, Eliopoulos GM, Sande MA. *The Sanford Guide to Antimicrobial Therapy*. 37th ed. Hyde Park, VT; 2007
- 87- van Kasteren ME, Manni n J, Ott A, Kullberg BJ, de Boer AS, Gyssens IC. Antibiotic prophylaxis and the risk of surgical site infections following total hip arthroplasty: timely administration is the most important factor. *Clin Infect Dis.* 2007; 44 (7): 921-7

- 88- Berghella V, Baxter JK, Chauhan SP. Evidence-based surgery for cesarean delivery. *Am J Obstet Gynecol.* 2005; 193 (5): 1607-17
- 89- Goldmann DA, Hopkins CC, Karchmer AW, Abel RM, McEnany MT, Akins C, Buckley MJ, Moellering RC Jr. Cephalothin prophylaxis in cardiac valve surgery. A prospective, double-blind comparison of two-day and six-day regimens. *J Thorac Cardiovasc Surg.* 1977; 73 (3): 470-9
- 90- Stone HH, Haney BB, Kolb LD, Geheber CE, Hooper CA. Prophylactic and preventive antibiotic therapy: timing, duration and economics. *Ann Surg.* 1979; 189 (6): 691-9
- 91- Pollard JP, Hughes SP, Scott JE, Evans MJ, Benson MK. Antibiotic prophylaxis in total hip replacement. *Br Med J.* 1979; 1 (6165): 707-9
- 92- Scher KS, Wroczynski AF, Jones CW. Duration of antibiotic prophylaxis. An experimental study. *Am J Surg.* 1986; 151 (2): 209-12
- 93- DiPiro JT, Cheung RP, Bowden TA Jr, Mansberger JA. Single dose systemic antibiotic prophylaxis of surgical wound infections. *Am J Surg.* 1986; 152 (5): 552-9
- 94- Rowe-Jones DC, Peel AL, Kingston RD, Shaw JF, Teasdale C, Cole DS. Single dose cefotaxime plus metronidazole versus three dose cefuroxime plus metronidazole as prophylaxis against wound infection in colorectal surgery: multicentre prospective randomised study. *BMJ.* 1990; 300 (6716): 18-22
- 95- Aberg C, Thore M. Single versus triple dose antimicrobial prophylaxis in elective abdominal surgery and the impact on bacterial ecology. *J Hosp Infect.* 1991; 18 (2): 149-54
- 96- Shapiro M. Prophylaxis in otolaryngologic surgery and neurosurgery: a critical review. *Rev Infect Dis.* 1991; 13 Suppl 10: S858-68
- 97- Boxma H, Broekhuizen T, Patka P, Oosting H. Randomised controlled trial of single-dose antibiotic prophylaxis in surgical treatment of closed fractures: the Dutch Trauma Trial. *Lancet.* 1996; 347 (9009): 1133-7
- 98- Scher KS. Studies on the duration of antibiotic administration for surgical prophylaxis. *Am Surg.* 1997; 63 (1): 59-62
- 99- Niederhäuser U, Vogt M, Vogt P, Genoni M, Kunzli A, Turina MI. Cardiac surgery in a high-risk group of patients: is prolonged postoperative antibiotic prophylaxis effective? *J Thorac Cardiovasc Surg.* 1997; 114(2): 162-8
- 100- Martin C, Pourriat JL. Quality of perioperative antibiotic administration by French anaesthetists. *J Hosp Infect.* 1998; 40 (1): 47-53
- 101- Glenny AM, Song F. Antimicrobial prophylaxis in colorectal surgery: a systematic review of randomized controlled trials. *Health Technol Assess.* 1998; 2 (7): 1-110

- 102- McDonald M, Grabsch E, Marshall C, Forbes A. Single-versus multiple-dose antimicrobial prophylaxis for major surgery: a systematic review. *Aust N Z J Surg.* 1998; 68 (6): 388-96
- 103- Namias N, Harvill S, Ball S, McKenney MG, Salomone JP, Civetta JM. Cost and morbidity associated with antibiotic prophylaxis in the ICU. *J Am Coli Surg.* 1999; 188 (3): 225-30
- 104- Coskun H, Erisen L, Basut O. Factors affecting wound infection rates in head and neck surgery. *Otolaryngol Head Neck Surg.* 2000; 123(3): 328-33
- 105- Yousuf M, Hussain M. Need and duration of antibiotic therapy in clean and clean contaminated operations. *J Pak Med Assoc.* 2002; 52 (7): 284-7
- 106- Velmahos GC, Toutouzas KG, Sarkisyan G, Chan LS, Jindal A, Karaiskakis M, Katkhouda N, Berne TV, Demetriades D. Severe trauma is not an excuse for prolonged antibiotic prophylaxis. *Arch Surg.* 2002; 137 (5): 537-41
- 107- Mui LM, Ng CS, Wong SK, Lam YH, Fung TM, Fok KL, Chung SS, Ng EK. Optimum duration of prophylactic antibiotics in acute non-perforated appendicitis. *ANI J Surg.* 2005; 75 (6): 425-8
- 108- Su HY, Ding DC, Chen DC, Lu MF, Liu JY, Chang FY. Prospective randomized comparison of single-dose versus 1-day cefazolin for prophylaxis in gynecologic surgery. *Ada Obstet Gynecol Scand.* 2005; 84 (4): 384-9
- 109- Takahashi S, Takeyama K, Miyamoto S, Tanuma Y, Takagi Y. Surgical antimicrobial prophylaxis in transurethral ureterolithotripsy. *J Infect Chemother.* 2005; 11 (5): 239-43
- 110- Fonseca SN, Kunzle SR, Junqueira MJ, Nascimento RT, de Andrade JL, Levin AS. Implementing 1-dose antibiotic prophylaxis for prevention of surgical site infection. *Arch Surg.* 2006; 141 (11): 1109-13
- 111- Avery CM, Ameerally P, Castling B, Swann RA. Infection of surgical wounds in the maxillofacial region and free flap donor sites with methicillin-resistant *Staphylococcus aureus*. *Br J Orai Maxillofac Surg.* 2006; 44 (3): 217-21
- 112- Andrews PJ, East CA, Jayaraj SM, Badia L, Panagamuwa C, Harding L. Prophylactic vs postoperative antibiotic use in complex septorhinoplasty surgery: a prospective, randomized, single-blind trial comparing efficacy. *Arch Facial Plast Surg.* 2006; 8 (2): 84-7
- 113- Liu SA, Tung KC, Shiao JY, Chiù YT. Preliminary report of associated factors in wound infection after major head and neck neoplasm operations-does the duration of prophylactic antibiotic matter? *Laryngol Otol.* 2008; 122 (4): 403-8
- 114- Nix DE, DiPiro JT, Bowden TA Jr, Vallner JJ. Cephalosporins for surgical prophylaxis: computer projections of intraoperative availability. *South Med J.* 1985; 78 (8): 962-6

- 115- DiPiro JT, Vallner JJ, Bowden TA Jr, Clark BA, Sisley JF
Intraoperative serum and tissue activity of cefazolin and cefoxitin. *Arch Surg.* 1985; 120 (7): 829-32
- 116- Levy M, Egersegi P, Strong A, Tessoro A, Spino M, Bannatyne R, Fear D, Posnick JC, Koren G. Pharmacokinetic analysis of cloxacillin loss in children undergoing major surgery with massive bleeding *Antimicrob Agents Chemother.* 1990; 34 (6): 1150-3
- 117- Wollinsky KH, Buchele M, Oethinger M, Kluger P, Mehrkens HH, Marre R, Puhl W. Influence of hemodilution on cefuroxime levels and bacterial contamination of intra- and postoperative processed wound blood during hip replacement. *Beitr Infusionsther Transfusionsmed.* 1996; 33: 191-5
- 118- Swoboda SM, Merz C, Kostuik J, Trentler B, Lipsett PA Does intraoperative blood loss affect antibiotic serum and tissue concentrations? *Arch Surg.* 1996; 131 (11): 1165-71
- 119- [No authors listed]. Antibiotic prophylaxis in surgery: summary of a Swedish-Norwegian Consensus Conference. *Scand J Infect Dis.* 1998; 30 (6): 547-57
- 120- Ohge H, Takesue Y, Yokoyama T, Murakami Y, Hiyama E, Yokoyama Y, Kanehiro T, Itaha H, Matsuura Y. An additional dose of cefazolin for intraoperative prophylaxis. *Surg Today.* 1999; 29(12): 1233-6
- 121- Zanetti G, Giardina R, Platt R. Intraoperative redosing of cefazolin and risk for surgical site infection in cardiac surgery. *Emerg Infect Dis.* 2001 ; 7 (5): 828-31
- 122- Dehne MG, Mühling J, Sablotzki A, Nopens H, Hempelmann G. Pharmacokinetics of antibiotic prophylaxis in major orthopedic surgery and blood-saving techniques. *Orthopedics.* 2001; 24 (7): 665-9
- 123- Gould FK, Elliott TS, Foweraker J, Fulford M, Perry JD, Roberts GJ, Sandoe JA, Watkin RW, Working Party of the British Society for Antimicrobial Chemotherapy. Guidelines for the prevention of endocarditis: report of the Working Party of the British Society for Antimicrobial Chemotherapy. *J Antimicrobial Chemother.* 2006; 57 (6): 1035-42
- 124- Wilson W, Taubert KA, Gewitz M, Lockhart PB, Baddour LM, Levison M, Bolger A, Cabell CH, Takahashi M, Baltimore RS, Newburger JW, Strom BL, Tani LY, Gerber M, Bonow RO, Pallasch T, Shulman ST, Rowley AH, Burns JC, Ferrieri P, Gardner T, Goff D, Durack DT; American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee; American Heart Association Council on Cardiovascular Disease in the Young; American Heart Association Council on Clinical Cardiology; American Heart Association Council on Cardiovascular Surgery and Anesthesia;

- Quality of Care and Outcomes Research Interdisciplinary Working Group
Prevention of infective endocarditis. Prevention of infective endocarditis: guidelines from the American Heart Association: a guideline from the American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee, Council on Cardiovascular Disease in the Young, and the Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and the Quality of Care and Outcomes Research Interdisciplinary Working Group *Circulation* 2007; 116 (15): 1736-54
- 125- Polk HC Jr, Trachtenberg L, Finn MP. Antibiotic activity in surgical incisions. The basis of prophylaxis in selected operations. *JAMA*. 1980; 244 (12): 1353-4
- 126- Waddell TK, Rotstein OD. Antimicrobial prophylaxis in surgery. Committee on Antimicrobial Agents, Canadian Infectious Disease Society. *CMAJ*. 1994; 151 (7): 925-31
- 127- Martin C. Antimicrobial prophylaxis in surgery: general concepts and clinical guidelines. French Study Group on Antimicrobial Prophylaxis in Surgery, French Society of Anesthesia and Intensive Care. *Infect Control Hosp Epidemiol*. 1994; 15(7): 463-71
- 128- Geroulanos S, Marathias K, Kriaras J, Kadas B. Cephalosporins in surgical prophylaxis. *J Chemother*. 2001; 13 Spec Noi (1): 23-6
- 129- Bolon MK, Morlote M, Weber SG, Koplan B, Carmeli Y, Wright SB. Glycopeptides are no more effective than beta-lactam agents for prevention of surgical site infection after cardiac surgery: a meta-analysis. *Clin Infect Dis*. 2004; 38 (10): 1357-63
- 130- [No authors listed]. National Nosocomial Infections Surveillance (NNIS) report, data summary from October 1986-April 1997, issued May 1997. A report from the NNIS System. *Am J Infect Control*. 1997; 25 (6): 477-87
- 131- Argenterò P.A, Campobasso G.A, Farina E.C, Vietti A.M. Linee guida per la prevenzione delle infezioni del sito chirurgico, 1999. Traduzione italiana di "Guideline for Prevention of Surgical Site Infection" Center for Disease Control, Atlanta, U.S.A., 1999. Regione Piemonte. Assessorato Sanità. www.regione.piemonte.it/sanita/sanpub/dwd/infezioni.pdf
- 132- Davis DA, Taylor-Vaisey A. Translating guidelines into practice. A systematic review of theoretic concepts, practical experience and research evidence in the adoption of clinical practice guidelines. *CMAJ*. 1997; 157(4): 408-16
- 133- Cabana MD, Rand CS, Powe NR, Wu AW, Wilson MH, Abboud PA, Rubin HR. Why don't physicians follow clinical practice guidelines? A framework for improvement. *JAMA*. 1999; 282(15): 1458-65

- 134- Burke JP. Maximizing appropriate antibiotic prophylaxis for surgical patients: an update from LDS Hospital, Salt Lake City. *Clin Infect Dis*. 2001; 33 Suppl 2: S78-83
- 135- Brusaferro S, Rinaldi O, Pea F, Faruzzo A, Barbone F. Protocol implementation in hospital infection control practice: an Italian experience of preoperative antibiotic prophylaxis. *J Hosp Infect*. 2001; 47 (4): 288-93
- 136- Gross PA, Pujat D. Implementing practice guidelines for appropriate antimicrobial usage: a systematic review. *Med Care*. 2001; 39 (8 Suppl 2): I155-69
- 137- Burnett KM, Scott MG, Kearney PM, Humphreys WG, McMillen RM. The identification of barriers preventing the successful implementation of a surgical prophylaxis protocol. *Pharm World Sci*. 2002; 24 (5): 182-7
- 138- Alerany C, Company D, Monterde J, Semeraro C. Impact of local guidelines and an integrated dispensing system on antibiotic prophylaxis quality in a surgical centre. *J Hosp Infect*. 2005; 60 (2): 111-7
- 139- Doebbeling BN, Pfaller MA, Kuhns KR, Massanari RM, Behrendt DM, Wenzel RP. Cardiovascular surgery prophylaxis. A randomized, controlled comparison of cefazolin and cefuroxime. *J Thorac Cardiovasc Surg*. 1990; 99 (6): 981-9
- 140- Townsend TR, Reitz BA, Bilker WB, Bartlett JG. Clinical trial of cefamandole, cefazolin, and cefuroxime for antibiotic prophylaxis in cardiac operations. *J Thorac Cardiovasc Surg*. 1993; 106(4): 664-70
- 141- Curtis JJ, Boley TM, Walls JT, Hamory B, Schmaltz RA. Randomized, prospective comparison of first- and second-generation cephalosporins as infection prophylaxis for cardiac surgery. *Am J Surg*. 1993; 166 (6): 734-7
- 142- Saginur R, Croteau D, Bergeron MG. Comparative efficacy of teicoplanin and cefazolin for cardiac operation prophylaxis in 3027 patients. The ESPRIT Group. *J Thorac Cardiovasc Surg*. 2000; 120 (6): 1120-30
- 143- Kriaras I, Michalopoulos A, Turina M, Geroulanos S. Evolution of antimicrobial prophylaxis in cardiovascular surgery, *Eur J Cardiothorac Surg*. 2000; 18 (4): 440-6
- 144- Bucknell SJ, Mohajeri M, Low J, McDonald M, Hill DG. Single- versus multiple-dose antibiotics prophylaxis for cardiac surgery. *Aust N Z J Surg*. 2000; 70 (6): 409-11
- 145- Finkelstein R, Rabino G, Mashiah T, Bar-El Y, Adler Z, Kertzman V, Cohen O, Milo S. Vancomycin versus cefazolin prophylaxis for cardiac surgery in the setting of a high prevalence of methicillin-resistant staphylococcal infections. *J Thorac Cardiovasc Surg*. 2002; 123 (2): 326-32

- 146- Bolon MK, Morlote M, Weber SG, Koplan B, Carmeli Y, Wright SB. Glycopeptides are no more effective than beta-lactam agents for prevention of surgical site infection after cardiac surgery: a meta-analysis. *Clin Infect Dis*. 2004; 38 (10): 1357-63
- 147- Miglioli PA, Merlo F, Calabro GB, Allerberger F, Fide M. Cefazolin concentrations in serum during cardiopulmonary bypass surgery. *Drugs Exp Clin Res*. 2005; 31 (1): 20-33
- 148- Dohmen PM. Influence of skin flora and preventive measures on surgical site infection during cardiac surgery. *Surg Infect (Larchmt)*. 2006; 7 Suppl 1: S13-7
- 149- Garey KW, Dao T, Chen H, Amrutkar P, Kumar N, Reiter M, Gentry LO. Timing of vancomycin prophylaxis for cardiac surgery patients and the risk of surgical site infections. *J Antimicrob Chemother*. 2006; 58 (3): 645-50
- 150- Edwards FH, Engelman RM, Houck P, Shahian DM, Bridges CR, Society of Thoracic Surgeons. The Society of Thoracic Surgeons Practice Guideline Series: Antibiotic Prophylaxis in Cardiac Surgery, Part I: Duration. *Ann Thorac Surg*. 2006; 81 (1): 397-404
- 151- Engelman R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, Jacobs M, Fernando H, Bridges C; Workforce on Evidence-Based Medicine, Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: Antibiotic prophylaxis in cardiac surgery, part II: Antibiotic choice. *Ann Thorac Surg*. 2007; 83 (4): 1569-76
- 152- Da Costa A, Kirkorian G, Cucherat M, Delahaye F, Chevalier P, Cerisier A, Isaaz K, Touboul P. Antibiotic prophylaxis for permanent pacemaker implantation: a meta-analysis. *Circulation*. 1998; 97(18): 1796-801
- 153- Bertaglia E, Zerbo F, Zardo S, Barzan D, Zoppo F, Pascotto P. Antibiotic prophylaxis with a single dose of cefazolin during pacemaker implantation: incidence of long-term infective complications. *Pacing Clin Electrophysiol*. 2006; 29 (1): 29-33
- 154- Sohail MR, Uslan DZ, Khan AH, Friedman PA, Hayes DL, Wilson WR, Steckelberg JM, Stoner SM, Baddour LM. Risk factor analysis of permanent pacemaker infection. *Clin Infect Dis*. 2007; 45 (2): 166-73
- 155- Klug D, Balde M, Pavin D, Hidden-Lucet F, Clementy J, Sadoul N, Rey JL, Lande G, Lazarus A, Victor J, Barnay C, Grandbastien B, Kacet S; PEOPLE Study Group. Risk factors related to infections of implanted pacemakers and cardioverter-defibrillators: results of a large prospective study. *Circulation*. 2007; 116 (12): 1349-55
- 156- Eagle KA, Guyton RA, Davidoff R, Edwards FH, Ewy GA, Gardner TJ, Hart JC, Herrmann HC, Hillis LD, Hutter AM Jr, Lytle BW, Marlow RA, Nugent WC, Orszulak TA, Antman EM, Smith

- SC Jr, Alpert JS, Anderson JL, Faxon DP, Fuster V, Gibbons RJ, Gregoratos G, Halperin JL, Hiratzka LF, Hunt SA, Jacobs AK, Ornato JP; American College of Cardiology; American Heart Association Task Force on Practice Guidelines; American Society for Thoracic Surgery and the Society of Thoracic Surgeons ACC/AHA 2004 guideline update for coronary artery bypass graft surgery: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1999 Guidelines for Coronary Artery Bypass Graft Surgery). *Circulation*. 2004; 110 (9): 1168-76
- 157- Pezzella AT, Adebonojo SA, Hooker SG, Mabogunje OA, Conlan AA. Complications of general thoracic surgery. *Curr Probi Surg*. 2000; 37(11): 733-858
- 158- Kaiser AB, Clayson KR, Mulherin JL Jr, Roach AC, Alien TR, Edwards WH, Dale WA. Antibiotic prophylaxis in vascular surgery. *Ann Surg*. 1978; 188 (3): 283-9
- 159- Pitt HA, Postier RG, MacGowan AW, Frank LW, Surmak AJ, Sitzman JV, Bouchier-Hayes D. Prophylactic antibiotics in vascular surgery. Topical, systemic, or both? *Ann Surg*. 1980; 192(3): 356-64
- 160- Edwards WH Jr, Kaiser AB, Kernodle DS, Appleby TC, Edwards WH Sr, Martin RS 3rd, Mulherin JL Jr, Wood CA Jr. Cefuroxime versus cefazolin as prophylaxis in vascular surgery. *J Vasc Surg*. 1992; 15(1): 35-41
- 161- Edwards WH Jr, Kaiser AB, Tapper S, Edwards WH Sr, Martin RS 3rd, Mulherin JL Jr, Jenkins JM, Roach AC. Cefamandole versus cefazolin in vascular surgical wound infection prophylaxis: cost-effectiveness and risk factors. *J Vasc Surg*. 1993; 18(3): 470-5
- 162- Bandyk DF. Antibiotics-why so many and when should we use them? *Semin Vasc Surg*. 2002; 15 (4): 268-74
- 163- Vardakas KZ, Soteriades ES, Chrysanthopoulou SA, Papagelopoulos PJ, Falagas ME. Perioperative anti-infective prophylaxis with teicoplanin compared to cephalosporins in orthopaedic and vascular surgery involving prosthetic material. *Clin Microbiol Infect*. 2005; 11 (10): 775-7
- 164- Stewart A, Eyers PS, Earnshaw JJ. Prevention of infection in arterial reconstruction. *Cochrane Database Syst Rev*. 2006; 3: CD003073
- 165- Stewart AH, Eyers PS, Earnshaw JJ. Prevention of infection in peripheral arterial reconstruction: a systematic review and meta-analysis. *J Vasc Surg*. 2007; 46 (1): 148-55
- 166- Società Oftalmologica Italiana. Raccomandazioni clinico organizzative per la chirurgia della cataratta non in regime di ricovero ordinario, 2000. (<http://www.sedesoi.com/>)

- 167- The Royal College of Ophthalmologists. Cataract surgery guidelines 2004. (<http://www.rcophth.ac.uk/>)
- 168- American Academy of Ophthalmology. Cataract in the Adult Eye, Preferred Practice Pattern. San Francisco: American Academy of Ophthalmology, 2006. (<http://one.aao.org/CE/PracticeGuidelines/PPP.aspx>)
- 169- Berkelman RL, Holland BW, Anderson RL. Increased bactericidal activity of dilute preparations of povidone-iodine solutions. *J Clin Microbiol.* 1982; 15 (4): 635-9
- 170- Apt L, Isenberg S, Yoshimori R, Paez JH. Chemical preparation of the eye in ophthalmic surgery. III. Effect of povidone-iodine on the conjunctiva. *Arch Ophthalmol.* 1984; 102 (5): 728-9
- 171- Speaker MG, Menikoff JA. Prophylaxis of endophthalmitis with topical povidone-iodine. *Ophthalmology* 1991; 98 (12): 1769-1775
- 172- Bohigian GM. A study of the incidence of culture-positive endophthalmitis after cataract surgery in an ambulatory care center. *Ophthalmic Surg Lasers.* 1999; 30 (4): 295-8
- 173- Ferguson AW, Scott JA, McGavigan J, Elton RA, McLean J, Schmidt U, Kelkar R, Dhillon B. Comparison of 5% povidone-iodine solution against 1% povidone-iodine solution in preoperative cataract surgery antisepsis: a prospective randomised double blind study. *Br J Ophthalmol.* 2003; 87 (2): 163-7
- 174- Mino de Kaspar H, Chang RT, Singh K, Egbert PR, Blumenkranz MS, Ta CN. Prospective randomized comparison of 2 different methods of 5% povidone-iodine applications for anterior segment intraocular surgery. *Arch Ophthalmol.* 2005; 123 (2): 161-5
- 175- Trinavarat A, Atchaneeyasakul LO, Nopmaneejumruslers C, Inson K. Reduction of endophthalmitis rate after cataract surgery with preoperative 5% povidone-iodine. *Dermatology.* 2006; 212 Suppl 1: 35-40
- 176- Ta CN, Singh K, Egbert PR, de Kaspar HM. Prospective comparative evaluation of povidone-iodine (10% for 5 minutes versus 5% for 1 minute) as prophylaxis for ophthalmic surgery. *J Cataract Refract Surg.* 2008; 34 (1): 171-2
- 177- Kaufman AH, Baum J. Penicillins and cephalosporins. In: Zimmerman TJ, ed, *Textbook of Ocular Pharmacology*. Philadelphia, Lippincott- Raven, 1997; 473-91
- 178- AAO-CDC Task Force on Vancomycin Prophylaxis in Ophthalmic Surgery. The Prophylactic Use of Vancomycin For Intraocular Surgery - Information Statement. American Academy of Ophthalmology, 1999.

<http://one.aao.org/CE/PracticeGuidelines/ClinicalStatements.aspx?sid=a3043761-ed 4-40a0-bb84-d353240d211 e>

- 179- Gordon YJ. Vancomycin prophylaxis and emerging resistance: are ophthalmologists the villains? The heroes? *Am J Ophthalmol.* 2001; 131 (3): 371-6
- 180- Wang RC, Lou PL, Ryan EA, Kroll AJ. Antibiotic therapy in post-operative endophthalmitis. *Semin Ophthalmol.* 2002; 17 (3-4): 153-61
- 181- Ciulla TA, Starr MB, Masket S. Bacterial endophthalmitis prophylaxis for cataract surgery: an evidence-based update. *Ophthalmology.* 2002; 109(1): 13-24
- 182- Libre PE, Della Latta P, Chin NX. Intracameral antibiotic agents for endophthalmitis prophylaxis: a pharmacokinetic model. *J Cataract Refract Surg.* 2003; 29(9): 1791-4
- 183- Benz MS, Scott IU, Flynn HW Jr, Unonius N, Miller D. Endophthalmitis isolates and antibiotic sensitivities: a 6-year review of culture-proven cases. *Am J Ophthalmology.* 2004; 137 (1): 38-42
- 184- Busbee BG. Advances in knowledge and treatment: an update on endophthalmitis. *Curr Opin Ophthalmol.* 2004; 15 (3): 232-7
- 185- Tipperman R. Pharmacologic considerations for cataract surgery. *Curr Opin Ophthalmol.* 2004; 15 (1): 51-5
- 186- Soriano ES, Nishi M. Endophthalmitis: incidence and prevention. *Curr Opin Ophthalmol.* 2005; 16 (1): 65-70
- 187- Wejde G, Samolov B, Seregard S, Koranyi G, Montan PG. Risk factors for endophthalmitis following cataract surgery: a retrospective case-control study. *J Hosp Infect.* 2005; 61 (3): 251-6
- 188- Garat M, Moser CL, Alonso-Tarrés C, Martin-Baranera M, Alberdi A. Intracameral cefazolin to prevent endophthalmitis in cataract surgery: 3-year retrospective study. *J Cataract Refract Surg.* 2005; 31 (11): 2230-4
- 189- Romero P, Méndez I, Salvat M, Fernández J, Almena M. Intracameral cefazolin as prophylaxis against endophthalmitis in cataract surgery, *J Cataract Refract Surg.* 2006; 32 (3): 438-41
- 190- Barry P, Seal DV, Gettinby G, Lees F, Peterson M, Revie CW; ESCRS Endophthalmitis Study Group. ESCRS study of prophylaxis of postoperative endophthalmitis after cataract surgery: Preliminary report of principal results from a European multicenter study. *J Cataract Refract Surg.* 2006; 32 (3): 407-10
- 191- Seal DV, Barry P, Gettinby G, Lees F, Peterson M, Revie CW, Wilhelmus KR; ESCRS Endophthalmitis Study Group. ESCRS study of prophylaxis of postoperative endophthalmitis after cataract surgery: Case for a European multicenter study. *J Cataract Refract Surg.* 2006; 32 (3): 396-406

- 192- Endophthalmitis Study Group, European Society of Cataract & Refractive Surgeons. Prophylaxis of postoperative endophthalmitis following cataract surgery: results of the ESCRS multicenter study and identification of risk factors. *J Cataract Refract Surg.* 2007; 33 (6): 978-88
- 193- Chang DF, Braga-Mele R, Mamalis N, Masket S, Miller KM, Nichamin LD, Packard RB, Packer M; ASCRS Cataract Clinical Committee. Prophylaxis of postoperative endophthalmitis after cataract surgery: results of the 2007 ASCRS member survey. *J Cataract Refract Surg.* 2007; 33 (10): 1801-5
- 194- Murphy CC, Nicholson S, Quah SA, Batterbury M, Neal T, Kaye SB. Pharmacokinetics of vancomycin following intracameral bolus injection in patients undergoing phacoemulsification cataract surgery. *Br J Ophthalmol.* 2007; 91 (10): 1350-3
- 195- Thoms SS, Musch DC, Soong HK. Postoperative endophthalmitis associated with sutured versus unsutured clear corneal cataract incisions. *Br J Ophthalmol.* 2007; 91 (6): 728-30
- 196- DeCroos FC, Afshari NA. Perioperative antibiotics and anti-inflammatory agents in cataract surgery. *Curr Opin Ophthalmol.* 2008; 19(1): 22-6
- 197- Mino de Kaspar H, Kreutzer TC, Aguirre-Romo I, Ta CN, Dudichum J, Bayrhof M, Klauss V, Kampik A. A prospective randomized study to determine the efficacy of preoperative topical levofloxacin in reducing conjunctival bacterial flora. *Am J Ophthalmol.* 2008; 145(1): 136-142
- 198- ASL 3 Regione Piemonte. "Protocolli di antibiotico profilassi in chirurgia O.R.L." Settembre 2002.
(<http://www.pri-asl3to.it/Protocolli/Schede/otorino.pdf>)
- 199- David N.F. Fairbanks. Pocket Guide to Antimicrobial Therapy in Otolaryngology - Head & Neck Surgery. The American Academy of Otolaryngology - Head and Neck Surgery Foundation, Inc, 13th Edition, 2007, pp. 68-72.
(<http://www.entnet.org/EducationAndResearch/upload/AAO-PGS-9-4-2.pdf>)
- 200- Becker GD, Pareli GJ. Cefazolin prophylaxis in head and neck cancer surgery. *Ann Otol Rhinol Laryngol.* 1979; 88 (2 Pt 1): 183-6
- 201- Righi M, Manfredi R, Farneti G, Pasquini E, Cenacchi V. Short-term versus long-term antimicrobial prophylaxis in oncologic head and neck surgery. *Head Neck.* 1996; 18 (5): 399-404
- 202- Rodrigo JP, Alvarez JC, Gómez JR, Suárez C, Fernández JA, Martfnez JA. Comparison of three prophylactic antibiotic regimen in clean-contaminated head and neck surgery. *Head Neck.* 1997; 19(3): 188-93

- 203- Carroll WR, Rosenstiel D, Fix JR, de la Torre J, Solomon JS, Brodish B, Rosenthal EL, Heinz T, Niwas S, Peters GE. Three-dose vs extended-course clindamycin prophylaxis for free flap reconstruction of the head and neck. *Arch Otolaryngol Head Neck Surg.* 2003; 129(7): 771-4
- 204- Shinagawa N, Yamanaka N, Suzuki K, Kawauchi H, Oda M, Yajin K, Ishizuka Y, Kurono Y, Iwai S, Yokoyama T, Takeyama H. A questionnaire survey on the theory of postoperative infection prophylaxis in otorhinolaryngology. *Jpn J Antibiot.* 2003; 56 (1): 15-26
- 205- Seven H, Sayin I, Turgut S. Antibiotic prophylaxis in clean neck dissections. *J Laryngol Otol.* 2004; 118 (3): 213-6
- 206- Ong SK, Morton RP, Kolbe J, Whitlock RM, McIvor NP. Pulmonary complications following major head and neck surgery with tracheostomy: a prospective, randomized, controlled trial of prophylactic antibiotics. *Arch Otolaryngol Head Neck Surg.* 2004; 130(9): 1084-7
- 207- Skitarelic N, Morovic M, Manestar D. Antibiotic prophylaxis in clean contaminated head and neck oncological surgery. *J Craniomaxillofac Surg.* 2007; 35 (1): 15-20
- 208- Lotfi CJ, Cavalcanti Rde C, Costa e Silva AM, Latorre Mdo R, Ribeiro Kde C, Carvalho AL, Kowalski LP. Risk factors for surgical-site infections in head and neck cancer surgery. *Otolaryngol Head Neck Surg.* 2008; 138 (1): 74-80
- 209- Verschuur HP, de Wever WW, van Benthem PP. Antibiotic prophylaxis in clean and clean-contaminated ear surgery. *Cochrane Database Syst Rev.* 2004; (3): CD003996
- 210- Annys E, Jorissen M. Short term effects of antibiotics (Zinnat) after endoscopic sinus surgery. *Acta Otorhinolaryngol Belg.* 2000; 54(1): 23-8
- 211- Rajan GP, Fergie N, Fischer U, Romer M, Radivojevic V, Hee GK. Antibiotic prophylaxis in septorhinoplasty? A prospective, randomized study. *Plast Reconstr Surg.* 2005; 116 (7): 1995-8
- 212- Kaygusuz I, Gök U, Yalgin S, Keles E, Kizirgil A, Demirbag E. Bacteremia during tonsillectomy. *Int J Pediatr Otorhinolaryngol.* 2001; 58(1): 69-73
- 213- Okur E, Arai M, Yildirim I, Kilic MA, Ciragil P. Bacteremia during adenoidectomy. *Int J Pediatr Otorhinolaryngol.* 2002; 66 (2): 149-53
- 214- Al-Kindy SA. Do antibiotics decrease post-tonsillectomy morbidity? *Saudi Med J.* 2002; 23 (6): 705-7
- 215- Piltcher OB, Scarton FB. Antibiotic use in tonsillectomies: therapeutic or prophylactic? Required or excessive? *Rev Bras Otorrinolaringol (Engl Ed).* 2005; 71 (5): 686-90

- 216- Burkart CM, Steward DL. Antibiotics for reduction of posttonsillectomy morbidity: a meta-analysis. *Laryngoscope*. 2005; 115(6): 997-1002
- 217- Sánchez-Carrión S, Prim MP, De Diego JI, Sastre N, Pena-Garcia P. Utility of prophylactic antibiotics in pediatric adenoidectomy. *Int J Pediatr Otorhinolaryngol*. 2006; 70 (7): 1275-81
- 218- Iyer S, DeFoor W, Grocela J, Kamholz K, Varughese A, Kenna M. The use of perioperative antibiotics in tonsillectomy: does it decrease morbidity? *Int J Pediatr Otorhinolaryngol*. 2006; 70 (5): 853-61
- 219- Dhiwakar M, Eng CY, Selvaraj S, McKerrow WS. Antibiotics to improve recovery following tonsillectomy: a systematic review. *Otolaryngol Head Neck Surg*. 2006; 134 (3): 357-64
- 220- Sistema Nazionale Linee Guida. SNLG 15 Appropriatezza e sicurezza degli interventi di tonsillectomia e/o adenoidectomia -Linea guida. Pubblicazione: Marzo 2008 (http://www.snlg-iss.it/cms/files/Snlq_Tonsille'08_interni.pdf)
- 221- Dhiwakar M, Clement W, Supriya M, McKerrow W. Antibiotics to reduce post-tonsillectomy morbidity. *Cochrane Database Syst Rev*. 2008; (2): CD005607
- 222- Gleisner AL, Argenta R, Pimentel M, Simon TK, Jungblut CF, Petteffi L, de Souza RM, Sauerssig M, Kruel CD, Machado AR. Infective complications according to duration of antibiotic treatment in acute abdomen. *Int J Infect Dis*. 2004, 8 (3): 155-62
- 223- Baum ML, Anish DS, Chalmers TC, Sacks HS, Smith H Jr, Fagerstrom RM. A survey of clinical trials of antibiotic prophylaxis in colon surgery: evidence against further use of no-treatment controls. *N Engl J Med*. 1981; 305 (14): 795-9
- 224- Price SA, Polk HC Jr. Prophylactic and therapeutic use of antibiotics in pelvic surgery. *J Surg Oncol*. 1999; 71 (4): 261-8
- 225- Mittelkötter U. Antimicrobial prophylaxis for abdominal surgery: is there a need for metronidazole? *J Chemother*. 2001; 13Spec No 1(1): 27-34
- 226- Piateli C, Hall JC. The prevention of wound infection in patients undergoing colorectal surgery. *J Hosp Infect*. 2001; 49 (4): 233-8
- 227- Rovera F, Diurni M, Dionigi G, Boni L, Ferrari A, Carcano G, Dionigi R. Antibiotic prophylaxis in colorectal surgery. *Expert Rev Anti Infect Ther*. 2005; 3 (5): 787-95
- 228- Koch A, Zippel R, Marusch F, Schmidt U, Gastinger I, Lippert H. Prospective multicenter study of antibiotic prophylaxis in operative treatment of appendicitis. *Dig Surg*. 2000; 17 (4): 370-8
- 229- Andersen BR, Kallehave FL, Andersen HK. Antibiotics versus placebo for prevention of postoperative infection after

- appendectomy. *Cochrane Database Syst Rev.* 2005; 20 (3): CD001439
- 230- Kasatpibal N, Norgaard M, Sorensen HT, Schonheyder HC, Jamulitrat S, Chongsuvivatwong V. Risk of surgical site infection and efficacy of antibiotic prophylaxis: a cohort study of appendectomy patients in Thailand. *BMC Infect Dis.* 2006; 6: 111
- 231- Al-Mulhim AS, Ali AM, Al-Masuod N, Alwahidi A. Post hemorrhoidectomy pain. A randomized controlled trial. *Saudi Med J.* 2006; 27(10): 1538-41
- 232- Wesarachawit W, Pattana-arun J. Antibiotics and early post operative complications of closed hemorrhoidectomy: a retrospective matched pair study. *J Med Assoc Thai.* 2007; 90 (9): 1828-32
- 233- Jewesson PJ, Stiver G, Wai A, Frighetto L, Nickoloff D, Smith J, Schwartz L, Sleigh K, Danforth D, Scudamore C, Chow A. Double-blind comparison of cefazolin and ceftizoxime for prophylaxis against infections following elective biliary tract surgery. *Antimicrob Agents Chemother.* 1996; 40 (1): 70-4
- 234- Westphal JF, Brogard JM. Biliary tract infections: a guide to drug treatment. *Drugs.* 1999; 57(1): 81-91
- 235- Illig KA, Schmidt E, Cavanaugh J, Krusch D, Sax HC. Are prophylactic antibiotics required for elective laparoscopic cholecystectomy? *J Am Coll Surg.* 1997; 184 (4): 353-6
- 236- Dobay KJ, Freier DT, Albear P. The absent role of prophylactic antibiotics in low-risk patients undergoing laparoscopic cholecystectomy. *Am Surg.* 1999; 65 (3): 226-8
- 237- McGuckin M, Shea JA, Schwartz JS. Infection and antimicrobial use in laparoscopic cholecystectomy. *Infect Control Hosp Epidemiol.* 1999; 20 (9): 624-6
- 238- Higgins A, London J, Charland S, Ratzner E, Clark J, Haun W, Maher DP. Prophylactic antibiotics for elective laparoscopic cholecystectomy: are they necessary? *Arch Surg.* 1999; 134 (6): 611-3
- 239- Tocchi A, Lepre L, Costa G, Liotta G, Mazzoni G, Maggiolini F. The need for antibiotic prophylaxis in elective laparoscopic cholecystectomy: a prospective randomized study. *Arch Surg.* 2000; 135(1): 67-70
- 240- Mahatharadol V. A reevaluation of antibiotic prophylaxis in laparoscopic cholecystectomy: a randomized controlled trial. *J Med Assoc Thai.* 2001; 84 (1): 105-8
- 241- Koc M, Zulfikaroglu B, Kece C, Ozalp N. A prospective randomized study of prophylactic antibiotics in elective laparoscopic cholecystectomy. *Surg Endosc.* 2003; 17 (11): 1716-8
- 242- Catarci M, Mancini S, Gentileschi P, Campione C, Sileri P, Grassi GB. Antibiotic prophylaxis in elective laparoscopic

- cholecystectomy. Lack of need or lack of evidence? *Surg Endosc.* 2004; 18(4): 638-41
- 243- Kuthe SA, Kaman L, Verma GR, Singh R. Evaluation of the role of prophylactic antibiotics in elective laparoscopic cholecystectomy: a prospective randomized trial. *Trop Gastroenterol.* 2006; 27 (1): 54-7
- 244- Chang WT, Lee KT, Chuang SC, Wang SN, Kuo KK, Chen JS, Sheen PC. The impact of prophylactic antibiotics on postoperative infection complication in elective laparoscopic cholecystectomy: a prospective randomized study. *Am J Surg.* 2006; 191 (6): 721-5
- 245- Platt R, Zucker JR, Zaleznik DF, Hopkins CC, Dellinger EP, Karchmer AW, Bryan CS, Burke JF, Wikler MA, Marino SK, et al. Perioperative antibiotic prophylaxis and wound infection following breast surgery. *J Antimicrob Chemother.* 1993; 31 Suppl B: 43-8
- 246- Hall JC, Hall JL. Antibiotic prophylaxis for patients undergoing breast surgery. *J Hosp Infect.* 2000; 46 (3): 165-70
- 247- D'Amico DF, Parimbelli P, Ruffolo C. Antibiotic prophylaxis in clean surgery: breast surgery and hernia repair. *J Chemother.* 2001; 13 Spec No 1(1):108-11
- 248- Tejirian T, DiFronzo LA, Haigh PI. Antibiotic prophylaxis for preventing wound infection after breast surgery: a systematic review and metaanalysis. *J Am Coli Surg.* 2006; 203 (5): 729-34
- 249- Cunningham M, Bunn F, Handscomb K. Prophylactic antibiotics to prevent surgical site infection after breast cancer surgery. *Cochrane Database Syst Rev.* 2006; 19 (2): CD005360
- 250- Gossner L, Keymling J, Hahn EG, EH C. Antibiotic prophylaxis in percutaneous endoscopic gastrostomy (PEG): a prospective randomized clinical trial. *Endoscopy.* 1999;31 (2): 119-24
- 251- Kulling D, Sonnenberg A, Fried M, Bauerfeind P. Cost analysis of antibiotic prophylaxis for PEG. *Gastrointest Endosc.* 2000; 51(2):152-6
- 252- Sharma VK, Howden CW. Meta-analysis of randomized, controlled trials of antibiotic prophylaxis before percutaneous endoscopic gastrostomy. *Am J Gastroenterol.* 2000; 95 (11): 3133-6
- 253- Nicholson FB, Korman MG, Richardson MA. Percutaneous endoscopic gastrostomy: a review of indications, complications and outcome. *J Gastroenterol Hepatol.* 2000; 15 (1): 21-5
- 254- Panigrahi H, Shreeve DR, Tan WC, Prudham R, Kaufman R. Role of antibiotic prophylaxis for wound infection in percutaneous endoscopic gastrostomy (PEG): result of a prospective double-blind randomized trial. *J Hosp Infect.* 2002; 50 (4): 312-5
- 255- Ahmad I, Mouncher A, Abdoolah A, Stenson R, Wright J, Daniels A, Tillett J, Hawthorne AB, Thomas G. Antibiotic

- prophylaxis for percutaneous endoscopic gastrostomy-a prospective, randomised, double-blind trial. *Aliment. Pharmacol Ther.* 2003; 18(2): 209-15
- 256- Saadeddin A, Freshwater DA, Fisher NC, Jones BJ. Antibiotic prophylaxis for percutaneous endoscopic gastrostomy for non-malignant conditions: a double-blind prospective randomized controlled trial. *Aliment. Pharmacol Ther.* 2005; 22 (6): 565-70
- 257- Lipp A, Lusardi G. Systemic antimicrobial prophylaxis for percutaneous endoscopic gastrostomy. *Cochrane Database Syst Rev.* 2006; 18 (4):CD005571
- 258- Imamura H, Furukawa H, Iijima S, Sugihara S, Tsujinaka T, Tsukuma H, Shimokawa T. Multicenter phase II study of antimicrobial prophylaxis in low-risk patients undergoing distal gastrectomy for gastric cancer. *Gastric Cancer.* 2006; 9(1): 32-5
- 259- Mohri Y, Tonouchi H, Kobayashi M, Nakai K, Kusunoki M, Mie Surgical Infection Research Group. Randomized clinical trial of single-versus multiple-dose antimicrobial prophylaxis in gastric cancer surgery. *Br J Surg.* 2007; 94 (6): 683-8
- 260- Rios A, Rodriguez JM, Munitiz V, Alcaraz P, Perez Flores D, Parrilla P. Antibiotic prophylaxis in incisional hernia repair using a prosthesis. *Hernia.* 2001; 5 (3): 148-52
- 261- Celdrà A, Frieyro O, de la Pinta JC, Souto JL, Esteban J, Rubio JM, Seharfs JF. The role of antibiotic prophylaxis on wound infection after mesh hernia repair under local anesthesia on an ambulatory basis. *Hernia.* 2004; 8 (1): 20-2
- 262- Aufenacker TJ, van Geldere D, van Mesdag T, Bossers AN, Dekker B, Scheijde E, van Nieuwenhuizen R, Hiemstra E, Maduro JH, Juttmann JW, Hofstede D, van Der Linden CT, Gouma DJ, Simons MP. The role of antibiotic prophylaxis in prevention of wound infection after Lichtenstein open mesh repair of primary inguinal hernia: a multicenter double-blind randomized controlled trial. *Ann Surg.* 2004; 240 (6): 955-60
- 263- Perez AR, Roxas MF, Hilvano SS. A randomized, double-blind, placebo-controlled trial to determine effectiveness of antibiotic prophylaxis for tension-free mesh herniorrhaphy. *J Am Coll Surg.* 2005; 200 (3): 393-7
- 264- Biswas S. Elective inguinal hernia repair with mesh: is there a need for antibiotic prophylaxis? - A review. *World J Surg.* 2005; 29 (7): 830-6
- 265- Aufenacker TJ, Koelemay MJ, Gouma DJ, Simons MP. Systematic review and meta analysis of the effectiveness of antibiotic prophylaxis in prevention of wound infection after mesh repair of abdominal wall hernia. *Br J Surg.* 2006; 93(1): 5-10

- 266- Terzi C. Antimicrobial prophylaxis in clean surgery with special focus on inguinal hernia repair with mesh. *J Hosp Infect.* 2006; 62 (4): 427-36
- 267- Tzovaras G, Delikoukos S, Christodoulides G, Spyridakis M, Mantzos F, Tepetes K, Athanassiou E, Hatzitheofilou C. The role of antibiotic prophylaxis in elective tension-free mesh inguinal hernia repair: results of a single-centre prospective randomised trial. *Int J Clin Pract.* 2007; 61 (2): 236-9
- 268- Sanabria A, Dominguez LC, Valdivieso E, Gómez G. Prophylactic antibiotics for mesh inguinal hernioplasty: a meta-analysis. *Ann Surg.* 2007; 245 (3): 392-6
- 269- Sanchez-Manuel FJ, Lozano-Garcia J, Seco-Gil JL. Antibiotic prophylaxis for hernia repair. *Cochrane Database Syst Rev.* 2007; (3): CD003769
- 270- Dorflinger T, Madsen PO. Antibiotic prophylaxis in transurethral surgery: a comparison of sulbactam-ampicillin and cefoxitin. *Infection.* 1985; 13 (2): 66-9
- 271- Botto H. Antibiotic prophylaxis in urology. Surgical and endoscopic surgery. Lithotripsy. Transplantation. *Ann Fr Anesth Reanim.* 1994; 13 (5 Suppl): S110-7
- 272- Llanes González L, Ruiz de la Roja JC, Martín Osés E, de Paz Cruz L, Zarate Rodríguez E, Sánchez Sánchez E, Paez Borda A, Llorente Abarca C, Berenguer Sánchez A. Antimicrobial prophylaxis in urology. *Actas Urol Esp.* 1997; 21 (6): 540-8
- 273- Kraklau DM, Wolf JS Jr. Review of antibiotic prophylaxis recommendations for office based urologic procedures. *Techn Urol.* 1999; 5: 123-128
- 274- Grabe M. Perioperative antibiotic prophylaxis in urology. *Curr Opinion Urol.* 2001 ; 11: 81 -85
- 275- Grabe M, Shortliffe L, Lobel B et al. Risk factors. In: Naber KG, Pechère JC, Kumazawa J et al., eds. Nosocomial and health care associated infections in urology. Health Publications Ltd, 2001, pp. 35-57
- 276- Olson ES, Cookson BD. Do antimicrobials have a role in preventing septicaemia following instrumentation of the urinary tract? *J Hosp Infect.* 2000; 45 (2): 85-97
- 277- Grabe M. Controversies in antibiotic prophylaxis in urology. *Int J Antimicrob Agents.* 2004; 23 Suppl 1 : S17-23
- 278- Takeyama K, Shimizu T, Mutoh M, Nishiyama N, Kunishima Y, Matsukawa M, Takahashi S, Hotta H, Itoh N, Tsukamoto T. Prophylactic antimicrobial agents in urologic laparoscopic surgery: 1-day versus 3-day treatments. *J Infect Chemother.* 2004; 10(3): 168-71
- 279- Johansen TE. Nosocomially acquired urinary tract infections in urology departments. Why an international prevalence study is

- needed in urology. *Int J Antimicrob Agents*. 2004; 23 Suppl 1 : S30-4
- 280- Matsumoto T, Kiyota H, Matsukawa M, Yasuda M, Arakawa S, Monden K; Japanese Society of UTI Cooperative Study Group (Chairman; Tetsuro Matsumoto). Japanese guidelines for prevention of perioperative infections in urological field. *Int J Urol*. 2007; 14(10): 890-909
- 281- Wolf JS Jr, Bennett CJ, Dmochowski RR, Hollenbeck BK, Pearle MS, Schaeffer AJ; Urologic Surgery Antimicrobial Prophylaxis Best Practice Policy Panel. Best practice policy statement on urologic surgery antimicrobial prophylaxis. *J Urol*. 2008; 179(4): 1379-90
- 282- Grabe M, Bishop M.C, Bjerklund-Johansen T.E, Botto H, Qek M, Lobel B, Naber K.G, Palou J, Tenke P. Guidelines on the management of urinary and male genital tract infections. ([http://www.uroweb.org/fileadmin/user_upload/Guidelines/The Management of Male Urinary and Genital Tract Infections.pdf](http://www.uroweb.org/fileadmin/user_upload/Guidelines/The%20Management%20of%20Male%20Urinary%20and%20Genital%20Tract%20Infections.pdf))
- 283- Yamamoto S, Shima H, Matsumoto T. Controversies in perioperative management and antimicrobial prophylaxis in urologic surgery. *Int J Urol*. 2008; 15 (6): 467-71
- 284- Taylor HM, Bingham JB. Antibiotic prophylaxis for transrectal prostate biopsy. *J Antimicrob Chemother*. 1997;39 (2): 115-117
- 285- Taylor HM, Bingham JB. The use of prophylactic antibiotics in ultrasound-guided transrectal prostate biopsy. *Clin Radiol*. 1997; 52 (10): 787-90
- 286- Sieber PR, Rommel FM, Agusta VE, Breslin JA, Huffnagle HW, Harpster LE. Antibiotic prophylaxis in ultrasound guided transrectal prostate biopsy. *J Urol*. 1997; 157 (6): 2199-200
- 287- Kapoor DA, Klimberg IW, Malek GH, Wegenke JD, Cox CE, Patterson AL, Graham E, Echols RM, Whalen E, Kowalsky SF. Single-dose oral ciprofloxacin versus placebo for prophylaxis during transrectal prostate biopsy. *Urology*. 1998; 52 (4): 552-8
- 288- Shandera KC, Thibault GP, Deshon GE Jr. Efficacy of one dose fluoroquinolone before prostate biopsy. *Urology*. 1998; 52 (4): 641-3
- 289- Isen K, Kupeli B, Sinik Z, Sözen S, Bozkirli I. Antibiotic prophylaxis for transrectal biopsy of the prostate: a prospective randomized study of the prophylactic use of single dose oral fluoroquinolone versus trimethoprim-sulfamethoxazole. *Int Urol Nephrol*. 1999; 31 (4): 491-5
- 290- Aron M, Rajeev TP, Gupta NP. Antibiotic prophylaxis for transrectal needle biopsy of the prostate: a randomized controlled study. *BJU Int*. 2000; 85 (6): 682-5
- 291- Griffith BC, Morey AF, Ali-Khan MM, Canby-Hagino E, Foley JP, Rozanski TA. Single dose levofloxacin prophylaxis for

- prostate biopsy in patients at low risk. *J Urol.* 2002; 168 (3): 1021-3
- 292- Webb NR, Woo HH. Antibiotic prophylaxis for prostate biopsy. *BJU Int.* 2002; 89: 824-828
- 293- Villers A, Mouton D, Rebillard X, Chautard D, Ruffion A, Staerman F, Comud F. Biopsy Technique and biopsy schemes for a first series of prostatic biopsies *Progr Urol.* 2004; 14 (2): 144-53
- 294- Sabbagh R, Me Cormack M, Peloquin F, Faucher R, Perreault JP, perrotte P, Karakiewicz PI, Saad F. A prospective randomized trial of 1-day versus 3-day antibiotic prophylaxis for transrectal ultrasound guided prostate biopsy. *Can J Urol.* 2004; 11 (2): 2216-9
- 295- Shigemura K, Tanaka K, Yasuda M, Ishihara S, Muratani T, Deguchi T, Matsumoto T, Kamidono S, Nakano Y, Arakawa S, Fujisawa M. Efficacy of 1-day prophylaxis medication with fluoroquinolone for prostate biopsy. *World J Urol.* 2005; 23 (5): 356-60
- 296- Raja J, Ramachandran N, Munneke G, Patel U. Current status of transrectal ultrasound-guided prostate biopsy in the diagnosis of prostate cancer. *Clin Radiol.* 2006; 61 (2): 142-53
- 297- Puig J, Darnell A, Bermúdez P, Malet A, Serrate G, Bare M, Prats J. Transrectal ultrasound-guided prostate biopsy: is antibiotic prophylaxis necessary? *Eur Radiol.* 2006; 16 (4): 939-43
- 298- Nielsen OS, Maigaard S, Frimodt-Moller N, Madsen PO. Prophylactic antibiotics in transurethral prostatectomy. *J Urol.* 1981; 126(1):60-2
- 299- Prokocimer P, Quazza M, Gibert C, Lemoine JE, Joly ML, Dureuil B, Moulouguet A, Manuel C, Desmots JM. Short-term prophylactic antibiotics in patients undergoing prostatectomy: report of a double-blind randomized trial with 2 intravenous doses of cefotaxime. *J Urol.* 1986; 135 (1): 60-4
- 300- Grabe M, Forsgren A, Bjork T, Hellsten S. Controlied trial of a short and a prolonged course with ciprofloxacin in patients undergoing transurethral prostatic surgery. *Eur J Clin Microbiol.* 1987; 6(1): 11-17
- 301- Grabe M. Antimicrobial agents in transurethral prostatic resection. *J Urol.* 1987; 138 (2): 245-52
- 302- Cox CE. Comparison of intravenous ciprofloxacin and intravenous cefotaxime for antimicrobial prophylaxis in transurethral surgery. *Am J Med.* 1989; 87 (5A): 252S-254S
- 303- Viitanen J, Talja M, Jussila E, Nurmi M, Permi J, Puolakka VM, Rintala E, salamela H, Tiitinen J, Tuhkanen K, et al. Randomized controlied study of chemoprophylaxis in transurethral prostatectomy. *J Urol.* 1993; 150 (5 Pt 2): 1715-7

- 304- Hargreave TB, Botto H, Rikken GH, Hindmarsh JR, McDermott TE, Mjolnerod OK, Petays P, Schalkhäuser K, Stellos A. European collaborative study of antibiotic prophylaxis for transurethral resection of the prostate. *Eur Urol.* 1993; 23 (4): 437-43
- 305- Lukkarinen O, Hellström P, Leppilähti M, Kontturi M, Tammela T. Antibiotic prophylaxis in patients with urinary retention undergoing transurethral prostatectomy. *Ann Ch'ir Gynaecol.* 1997; 86 (3): 239-42
- 306- Liu GG, Nguyen T, Nichol MB. An economic analysis of antimicrobial prophylaxis against urinary tract infection in patients undergoing transurethral resection of the prostate. *Clin Ther.* 1999; 21 (9): 1589-604
- 307- Berry A, Barratt A. Prophylactic antibiotic use in transurethral prostatic resection: a meta-analysis. *J Urol.* 2002; 167 (2 Pt 1): 571-7
- 308- Qiang W, Jianchen W, MacDonald R, Monga M, Wilt TJ. Antibiotic prophylaxis for transurethral prostatic resection in men with preoperative urine containing less than 100,000 bacteria per ml: a systematic review. *J Urol.* 2005; 173 (4): 1175-81
- 309- Wagenlehner FM, Wagenlehner C, Schinzel S, Naber KG; Working Group "Urological Infections" of German Society of Urology. Prospective, randomized, multicentric, open, comparative study on the efficacy of a prophylactic single dose of 500 mg levofloxacin versus 1920 mg trimethoprim/ sulfamethoxazole versus a control group in patients undergoing TUR of the prostate. *Eur Urol.* 2005; 47: 549-56
- 310- Terai A, Ichioka K, Kohei N, Ueda N, Utsunomiya N, Inoue K. Antibiotic prophylaxis in radical prostatectomy: 1-day versus 4-day treatments. *Int J Urol.* 2006; 13 (12): 1488-93
- 311- Takeyama K, Takahashi S, Maeda T, Mutoh M, Kunishima Y, Matsukawa M, Takagi Y. Comparison of 1-day, 2-day, and 3-day administration of antimicrobial prophylaxis in radical prostatectomy. *J Infect Chemother.* 2007; 13 (5): 320-3
- 312- Ozturk M, Koca O, Kaya C, Karaman MI. A prospective randomized and placebo-controlled study for the evaluation of antibiotic prophylaxis in transurethral resection of the prostate. *Urol Int.* 2007; 79(1): 37-40
- 313- Sakura M, Kawakami S, Yoshida S, Masuda H, Kobayashi T, Kihara K. Prospective comparative study of single dose versus 3-day administration of antimicrobial prophylaxis in minimum incision endoscopic radical prostatectomy. *Int J Urol.* 2008; 15 (4): 328-31
- 314- Charton M, Vallancien G, Veillon B, Brisset JM. Urinary tract infection in percutaneous surgery for renal calculi. *J Urol.* 1986; 135(1): 15-7

- 315- Pearle MS, Roehrborn CG. Antimicrobial prophylaxis prior to shock wave lithotripsy in patients with sterile urine before treatment: a meta-analysis and cost-effectiveness analysis. *Urology*. 1997; 49 (5): 679-86
- 316- Dogan HS, Sahin A, Cetinkaya Y, Akdogan B, Ozden E, Kendi S. Antibiotic prophylaxis in percutaneous nephrolithotomy: prospective study in 81 patients. *J Endourol*. 2002; 16 (9): 649-53
- 317- Knopf HJ, Graff HJ, Schulze H. Perioperative antibiotic prophylaxis in ureteroscopic stone removal. *Eur Urol*. 2003; 44 (1): 115-8
- 318- MacDermott JP, Ewing RE, Somerville JF, Gray BK. Cephadrine prophylaxis in transurethral procedures for carcinoma of the bladder. *Br J Urol*. 1988; 62 (2): 136-9
- 319- Wilson L, Ryan J, Thelning C, Masters J, Tuckey J. Is antibiotic prophylaxis required for flexible cystoscopy? A truncated randomized double-blind controlled trial. *J Endourol*. 2005; 19(8): 1006-8
- 320- Hara N, Kitamura Y, Saito T, Komatsubara S, Nishiyama T, Takahashi K. Perioperative antibiotics in radical cystectomy with ileal conduit urinary diversion: Efficacy and risk of antimicrobial prophylaxis on the operation day alone. *Int J Urol*. 2008; 15 (6): 511-5
- 321- McClelland S 3rd, Hall WA. Postoperative central nervous system infection: incidence and associated factors in 2111 neurosurgical procedures. *Clin Infect Dis*. 2007 45 (1): 55-9
- 322- Valentini LG, Casali C, Chatenoud L, Chiaffarino F, Uberti-Foppa C, Broggi G. Surgical site infections after elective neurosurgery: a survey of 1747 patients. *Neurosurgery*. 2008; 62 (1): 88-95
- 323- Langley JM, LeBlanc JC, Drake J, Milner R. Efficacy of antimicrobial prophylaxis in placement of cerebrospinal fluid shunts: meta-analysis. *Clin Infect Dis*. 1993;17 (1): 98-103
- 324- Haines SJ, Walters BC. Antibiotic prophylaxis for cerebrospinal fluid shunts: a metanalysis. *Neurosurgery*. 1994 34 (1): 87-92
- 325- Barker FG 2nd. Efficacy of prophylactic antibiotics for craniotomy: a meta-analysis. *Neurosurgery*. 1994; 35 (3): 484-90
- 326- Korinek AM, Golmard JL, Elcheick A, Bismuth R, van Effenterre R, Coriat P, Puybasset L. Risk factors for neurosurgical site infections after craniotomy: a critical appraisal of antibiotic prophylaxis on 4,578 patients. *Br J Neurosurg*. 2005; 19 (2): 155-62
- 327- Brown EM. Antibiotic prophylaxis in patients undergoing clean, non-implant craniotomy. *Br J Neurosurg*. 2006; 20 (5): 273-4

- 328- Ratilal B, Costa J, Sampaio C. Antibiotic prophylaxis for surgical introduction of intracranial ventricular shunts. *Cochrane Database Syst Rev.* 2006 19; 3: CD005365
- 329- Barker FG 2nd. Efficacy of prophylactic antibiotics against meningitis after craniotomy: a meta-analysis. *Neurosurgery.* 2007; 60 (5): 887-94
- 330- Brown SM, Anand VK, Tabaei A, Schwartz TH. Role of perioperative antibiotics in endoscopic skull base surgery. *Laryngoscope.* 2007; 117(9): 1528-32
- 331- Ratilal B, Costa J, Sampaio C. Antibiotic prophylaxis for surgical introduction of intracranial ventricular shunts: a systematic review. *J Neurosurg Pediatrics.* 2008; 1 (1): 48-56
- 332- Price SA, Polk HC Jr. Prophylactic and therapeutic use of antibiotics in pelvic surgery. *J Surg Oncol.* 1999; 71 (4): 261-8
- 333- Giuliani B, Periti E, Mecacci F. Antimicrobial prophylaxis in obstetric and gynecological surgery. *J Chemother.* 1999; 11 (6): 577-80
- 334- Guaschino S, De Santo D, De Seta F. New perspectives in antibiotic prophylaxis for obstetric and gynaecological surgery. *J Hosp Infect.* 2002; 50 Suppl A: S13-6
- 335- Cormio G, Di Fazio F, Lorusso F, Di Gesù G, Cacciapuoti C, Loverro G, Nappi L, Selvaggi L. Antimicrobial prophylaxis in laparotomie gynecologic surgery: a prospective randomized study comparing amoxicillin-clavulanic acid with cefazolin. *J Chemother.* 2002; 14 (6): 618-22
- 336- Larsen JW, Hager WD, Livengood CH, Hoyme U. Guidelines for the diagnosis, treatment and prevention of postoperative infections. *Infect Dis Obstet Gynecol.* 2003; 11(1): 65-70
- 337- Cormio G, Di Fazio F, Cacciapuoti C, Bettocchi S, Borraccino L, Selvaggi L. Prospective randomized study comparing amoxicillin-clavulanic acid with cefazolin as antimicrobial prophylaxis in laparotomie gynecologic surgery. *Acta Obstet Gynecol Scand.* 2003; 82 (12): 1130-4
- 338- Cormio G, Bettocchi S, Ceci O, Nappi L, Di Fazio F, Cacciapuoti C, Selvaggi L. Antimicrobial prophylaxis in laparoscopic gynecologic surgery: a prospective randomized study comparing amoxicillin-clavulanic acid with cefazolin. *J Chemother.* 2003; 15 (6): 574-8
- 339- Scardigno D, Lavermicocca T, Magazino MP, Cazzolla A, Lorusso M, Vicino M. A randomized study comparing amoxicillin-clavulanic acid with cefazolin as antimicrobial prophylaxis in laparotomie gynecologic surgery. *Minerva Ginecol.* 2006; 58 (2): 85-90
- 340- ACOG Committee on Practice Bulletins. ACOG Practice Bulletin No. 74. Antibiotic prophylaxis for gynecologic procedures. *Obstet Gynecol.* 2006; 108 (1): 225-34

- 341- Ledger WJ. Prophylactic antibiotics in obstetrics-gynecology: a current asset, a future liability? *Expert Rev Anti Infect Ther.* 2006; 4 (6): 957-64
- 342- Cormio G, Vicino M, Loizzi V, Tangari D, Selvaggi L. Antimicrobial prophylaxis in vaginal gynecologic surgery: a prospective randomized study comparing amoxicillin-clavulanic acid with cefazolin. *J Chemother.* 2007; 19 (2): 193-7
- 343- Popovic J, Grujic Z, Sabo A. Influence of pregnancy on ceftriaxone, cefazolin and gentamicin pharmacokinetics in caesarean vs. non-pregnant sectioned women. *J Clin Pharm Ther.* 2007; 32 (6): 595-602
- 344- Faro S, Martens MG, Hammill HA, Riddle G, Tortolero G. Antibiotic prophylaxis: is there a difference? *Am J Obstet Gynecol.* 1990; 162 (4): 900-7
- 345- Carlson C, Duff P. Antibiotic prophylaxis for cesarean delivery: is an extended-spectrum agent necessary? *Obstet Gynecol.* 1990; 76 (3 Pt 1): 343-6
- 346- Currier JS, Tosteson TD, Platt R. Cefazolin compared with cefoxitin for cesarean section prophylaxis: the use of a two-stage study design. *J Clin Epidemiol.* 1993; 46 (7): 625-30
- 347- Noyes N, Berkeley AS, Freedman K, Ledger W. Incidence of postpartum endomyometritis following single-dose antibiotic prophylaxis with either ampicillin/sulbactam, cefazolin, or cefotetan in high-risk cesarean section patients. *Infect Dis Obstet Gynecol.* 1998; 6(5): 220-3
- 348- Hopkins L, Smaill. Antibiotic prophylaxis regimens and drugs for cesarean section. *Cochrane Database Syst Rev.* 2000; (2): CD001136
- 349- Chelmow D, Rueli MS, Huang E. Prophylactic use of antibiotics for nonlaboring patients undergoing cesarean delivery with intact membranes: a meta-analysis. *Am J Obstet Gynecol.* 2001; 184 (4): 656-61
- 350- Killian CA, Graffunder EM, Vinciguerra TJ, Venezia RA. Risk factors for surgical-site infections following cesarean section. *Infect Control Hosp Epidemiol.* 2001 ; 22 (10): 613-7
- 351- Smaill F, Hofmeyr GJ. Antibiotic prophylaxis for cesarean section. *Cochrane Database Syst Rev.* 2002; (3): CD000933
- 352- American College of Obstetricians and Gynecologists. ACOG practice bulletin number 47, October 2003: Prophylactic Antibiotics in Labor and Delivery. *Obsfef Gynecol.* 2003; 102 (4): 875-82
- 353- Chelmow D, Hennesy M, Evantash EG. Prophylactic antibiotics for non-laboring patients with intact membranes undergoing cesarean delivery: an economic analysis. *Am J Obstet Gynecol.* 2004; 191 (5): 1661-5

- 354- Berghella V, Baxter JK, Chauhan SP. Evidence-based surgery for cesarean delivery. *Am J Obstet Gynecol.* 2005; 193 (5): 1607-17
- 355- Martins AC, Krauss-Silva L. Systematic reviews of antibiotic prophylaxis in cesareans. *Cad Saude Publica.* 2006; 22 (12): 2513-26
- 356- Olsen MA, Butler AM, Willers DM, Devkota P, Gross GA, Fraser VJ. Risk factors for surgical site infection after low transverse cesarean section. *Infect Control Hosp Epidemiol.* 2008; 29 (6): 477-84
- 357- Mittendorf R, Aronson MP, Berry RE, Williams MA, Kupelnick B, Klickstein A, Herbst AL, Chalmers TC. Avoiding serious infections associated with abdominal hysterectomy: a meta-analysis of antibiotic prophylaxis. *Am J Obstet Gynecol.* 1993; 169(5): 1119-24
- 358- Tanos V, Rojansky N. Prophylactic antibiotics in abdominal hysterectomy. *J Am Coli Surg.* 1994; 179 (5): 593-600
- 359- Meltomaa SS, Mäkinen JI, Taalikka MO, Helenius HY. Incidence, risk factors and outcome of infection in a 1-year hysterectomy cohort: a prospective follow-up study. *J Hosp Infect.* 2000; 45 (3): 211-7
- 360- Chongsomchai C, Lumbiganon P, Thinkhamrop J, Ounchai J, Vudhikamraksa N. Placebo-controlled, double-blind, randomized study of prophylactic antibiotics in elective abdominal hysterectomy. *J Hosp Infect.* 2002; 52 (4): 302-6
- 361- Löfgren M, Poromaa IS, Stjerndahl JH, Renström B. Postoperative infections and antibiotic prophylaxis for hysterectomy in Sweden: a study by the Swedish National Register for Gynecologic Surgery. *Ada Obstet Gynecol Scand.* 2004; 83(12): 1202-7
- 362- Eason EL, Wells GA, Garber GE, Hopkins ML; Vaginal Antisepsis for Abdominal Hysterectomy Study Group. Prophylactic antibiotics for abdominal hysterectomy: indication for low-risk Canadian women. *J Obstet Gynaecol Can.* 2004; 26 (12): 1067-72
- 363- Costa RJ, Krauss-Silva L. Systematic review and meta-analysis of antibiotic prophylaxis in abdominal hysterectomy. *Cad Saude Publica.* 2004; 20 Suppl 2: S175-89
- 364- Kocak I, Ustun C, Gürkan N. Prophylactic antibiotics in elective abdominal hysterectomy. *Int J Gynaecol Obstet.* 2005; 90 (2): 157-8
- 365- Chang WC, Hung YC, Li TC, Yang TC, Chen HY, Lin CC. Short course of prophylactic antibiotics in laparoscopically assisted vaginal hysterectomy. *J Reprod Med.* 2005; 50 (7): 524-8

- 366- Sawaya GF, Grady D, Kerlikowske K, Grimes DA. Antibiotics at the time of induced abortion: the case for universal prophylaxis based on a meta-analysis. *Obstet Gynecol.* 1996; 87 (5 Pt 2): 884-90
- 367- Penney GC. Preventing infective sequelae of abortion. *Hum Reprod.* 1997; 12(11 Suppl):107-12
- 368- Keder LM. Best practices in surgical abortion. *Am J Obstet Gynecol.* 2003;189 (2): 418-22
- 369- No authors listed. Antimicrobial prophylaxis for orthopaedic surgery. *Drug Ther Bull.* 2001 ; 39 (6): 43-6
- 370- Vardakas KZ, Soteriades ES, Chrysanthopoulou SA, Papagelopoulos PJ, Falagas ME. Perioperative anti-infective prophylaxis with teicoplanin compared to cephalosporins in orthopaedic and vascular surgery involving prosthetic material. *Clin Microbiol Infect.* 2005; 11 (10): 775-7
- 371- W-Dahl A, Toksvig-Larsen S. Infection prophylaxis: a prospective study in 106 patients operated on by tibial osteotomy using the hemicallotaxis technique. *Arch Orthop Trauma Surg.* 2006; 126 (7): 441-7
- 372- Kato D, Maezawa K, Yonezawa I, Iwase Y, Ikeda H, Nozawa M, Kurosawa H. Randomized prospective study on prophylactic antibiotics in clean orthopedic surgery in one ward for 1 year. *J Orthop Sci.* 2006; 11 (1): 20-7
- 373- Ali M, Raza A. Role of single dose antibiotic prophylaxis in clean orthopedic surgery. *Coli Physicians Surg Pak.* 2006; 16 (1): 45-8
- 374- Trampuz A, Zimmerli W. Antimicrobial agents in orthopaedic surgery: Prophylaxis and treatment. *Drugs.* 2006; 66 (8): 1089-105
- 375- Fletcher N, Sofianos D, Berkes MB, Obremskey WT. Prevention of perioperative infection. *J Bone Joint Surg Am.* 2007; 89(7): 1605-18
- 376- Prokuski L. Prophylactic antibiotics in orthopaedic surgery. *J Am Acad Orthop Surg.* 2008; 16 (5): 283-93
- 377- Glenny A, Song F. Antimicrobial prophylaxis in total hip replacement: a systematic review. *Health Technol Assess.* 1999; 3(21): 1-57
- 378- Hanssen AD, Osmon DR. The use of prophylactic antimicrobial agents during and after hip arthroplasty. *Clin Orthop RelatRes.* 1999; (369): 124-38
- 379- Eshkenazi AU, Garti A, Tamir L, Hendel D. Serum and synovial vancomycin concentrations following prophylactic administration in knee arthroplasty. *Am J Knee Surg.* 2001 Fall; 14(4): 221-3

- 380- Tang WM, Chiù KY, Ng TP, Yau WP, Ching PT, Seto WH. Efficacy of a single dose of cefazolin as a prophylactic antibiotic in primary arthroplasty. *J Arthroplasty*. 2003; 18 (6): 714-8
- 381- Gillespie WJ; Walenkamp G. Antibiotic Prophylaxis for surgery for proximal femoral and other closed long bone fractures. *Cochrane Database Syst Rev*. 2001; (1): CD000244
- 382- Gosselin RA, Roberts I, Gillespie WJ. Antibiotics for preventing infection in open limb fractures. *Cochrane Database Syst Rev*. 2004; (1): CD003764
- 383- Southwell-Keely JP, Russo RR, March L, Cumming R, Cameron I, Brnabic AJ. Antibiotic prophylaxis in hip fracture surgery: a metaanalysis. *Clin Orthop Relat Res*. 2004; (419): 179-84
- 384- Hauser CJ, Adams CA Jr, Eachempati SR; Council of the Surgical Infection Society. Surgical Infection Society guideline: prophylactic antibiotic use in open fractures: an evidence-based guideline. *Surg Infect (Larchmt)*. 2006; 7 (4): 379-405
- 385- Slobogean GP, Kennedy SA, Davidson D, O'Brien PJ. Single- Versus Multiple-Dose Antibiotic Prophylaxis in the Surgical Treatment of Closed Fractures: A Meta-Analysis. *J Orthop Trauma*. 2008; 22 (4): 264-269
- 386- Dimick JB, Lipsett PA, Kostuik JP. Spine update: antimicrobial prophylaxis in spine surgery: basic principles and recent advances. *Spine*. 2000; 25 (19): 2544-8
- 387- Barker FG 2nd. Efficacy of prophylactic antibiotic therapy in spinal surgery: a meta-analysis. *Neurosurgery*. 2002; 51 (2): 391-400
- 388- Mathis JM. Percutaneous vertebroplasty: complication avoidance and technique optimization. *AJNR Am J Neuroradiol*. 2003; 24(8): 1697-706
- 389- Dobzyniak MA, Fischgrund JS, Hankins S, Herkowitz HN. Ingle versus multiple dose antibiotic prophylaxis in lumbar disc surgery. *Spine*. 2003; 28 (21): E453-5
- 390- Brown EM, Pople IK, de Louvois J, Hedges A, Bayston R, Eisenstein SM, Lees P; British Society for Antimicrobial Chemotherapy Working Party on Neurosurgical Infections. Spine update: prevention of postoperative infection in patients undergoing spinal surgery. *Spine*. 2004; 29 (8): 938-45
- 391- Mastronardi L, Rychlicki F, Tatta C, Morabito L, Agrillo U, Ducati A. Spondylodiscitis after lumbar microdiscectomy: effectiveness of two protocols of intraoperative antibiotic prophylaxis in 1167 cases. *Neurosurg Rev*. 2005; 28 (4): 303-7
- 392- Kanayama M, Hashimoto T, Shigenobu K, Oha F, Togawa D. Effective prevention of surgical site infection using a Centers for Disease Control and Prevention guideline-based antimicrobial

